

THE OROANTRAL FISTULA: A CASE REPORT

^[1] Dr.M.veeramuthu, MDS, ^[2]Mrs. Sunitha MSC(N), ^[3]Dr. A. Mathan Mohan MDS, ^[4]Dr.GR Karthikeyan MDS, ^[5]Dr.V.Deepak MDS, ^[6]Dr.Nirmala devar MDS

^[1] Senior lecturer, Department of Oral and Maxillo Facial Surgery Karpaga Vinayaka Institute of Dental Sciences.

^[2] Associate professor Karpaka Vinayaka College of Nursing.

^[3] Professor, Department of oral and maxilla facial surgery Karpaka vinayaka institute of dental sciences

^[4] Reader, Department of oral and maxilla facial surgery Karpaka vinayaka institute of dental sciences

^[5] Reader, Department of oral and maxilla facial surgery Karpaka vinayaka institute of dental sciences

^[6] Senior lecturer, Department of Oral and Faxilla Facial surgery Karpaka Vinayaka Institute of Dental Sciences

ABSTRACT

Introduction: An unnatural communication between the maxillary sinus and oral cavity is known as “oro-antral communication”(OAC) and if it does not close spontaneously, it is epithelialized so that oroantral fistula develops.(2) The most common cause of oroantral fistula is the extraction of a maxillary molar or premolar.(2,11) The close relationship between the apex of these teeth and the thinness of the antral floor explains this.(11) From simple local methods like buccal advancement flap to complex distal flaps and grafts, various methods have been described in the literature for the closure of these communication.(2) To facilitate gravitational drainage and aeration via an inferior meatal antrostomy and to remove irreversibly damaged mucosa of the maxillary sinus, Caldwell Luc procedure was designed.(10) **Objective:** To describe a case report of oroantral fistula [OAF] and its closure. **Case Report:** We report a case of 28 years old male patient who came to our attention for the presence of the OAF and presence of maxillary third molar root piece in the maxillary sinus. This patient was treated by“ OBTURATOR procedure.”

Conclusion: This treatment modality provide a systematic approach for repair of oro-antral communications

INTRODUCTION

The existence of an unnatural communication between the oral cavity and maxillary sinus due to loss of soft and hard tissues that normally separate these compartments, characterizes the “oroantral communications”(OAC).(13) The oroantral fistula(OAF) is a pathological communication between the maxillary sinus and oral cavity and it can be classified as alveolosinus, palatal-sinus and vestibulo- sinus, depending on the location. (8) Dental infection, radiation therapy, sequelae of removal of maxillary cysts(10-15%) and tumors (5-10%), osteomyelitis, trauma(2-5%) can cause OAC.(6) The most common etiologic factor for oroantral communications(OAC) is upper molars extractions(0.31% - 4.7%).(12) Postop frequency of OAC varies between 3.8%(ARRIGONI &

LAMBRECHJ 2004) and 18.7% (ROTHAMEL ET AL. 2007).(7) Oroantral communication (OAC) is the most common complication in the maxilla due to the close proximity of the third molars to the maxillary sinus. From simple local methods like buccal advancement flap to complex distal flaps and grafts, various methods have been described in the literature for closure of these communication. (2) The Caldwell-Luc operation was first described while creating intranasal counter drainage through the inferior meatus in the late 19th century.(10) The Caldwell Luc operation was first described as a technique to remove infection and diseased mucosa from the maxillary sinus via the canine fossa.(10) The objective of this article to describe a case report of oroantral fistula and its management by non surgical method by fixed obturator with an extension closing the communication and allowing it heal by secondary

intention

Case Report A 48 year old Female patient had referred to the department of the “Oral and Maxillofacial Surgery”, karpaga vinayaka institute of dental sciences, with chief complaints of the pain in the upper left back teeth region and pain with the left buccal mucosa. Radiographic examination revealed the presence of the left maxillary third molar root piece in the maxillary sinus and presence of the oroantral communication. Fig. 1: OPG Report Showing the location of displaced root piece Under local anaesthesia administration, forighn obstructing the extracting removed epithelial lining removed and obturator placed with in extention in to the extracting socket the obturator is stabilise by using tie wire between 13,14 and 23,24 region . The patient’s follow-up was done carefully and was examined for the presence of oro-antral communication during follow-up



its relationship to the adjacent teeth, height of the alveolar ridge, persistence and sinus inflammation and the patient’s general health while choosing the surgical approach for treatment of an oroantral fistula.(8) A systematic antifungal treatment must be combined with abundant washing with saline and topical antifungal solution, because a chronic communication between the maxillary sinus and oral cavity can denote an access route for fungal penetration into the sinus.(8) required prior to antrum exposure to remove the diseased lining if severe sinusitis is present.(2) The success rate of

the primary closing of oro-antral fistulas in 48 hours is about 90-95% and it falls to 67% in secondary closing.(1) To close oro-antral communications, the buccal fat pad can be used.(3) Buccal fat pad do not interfere the vestibular groove depth and because of its anatomical position, it can be used as pedicellate graft, hence, buccal fat pad is more beneficial to treat oro-antral communications.(3) For the closure of OAF, Bio-Oss-Bio-Gide Sandwich technique has been used. This procedure is beneficial for achieving bony and soft tissue closure.(3) Allografts, autografts, absorbable materials [e.g: polydioxanon], synthetic materials [e.g. gold foils], rotational flaps are the different treatment modalities for the treatment of Oroantral fistula.(11)



Discussion Maxillary sinus is a part of the paranasal

sinuses which is internally revested by a membrane known as “schneiderian membrane”.(11) “Highmore antrum” is the another name of the maxillary sinus. (11) The most important paranasal sinuses are the maxillary sinuses because of their proximity to the roots of maxillary dentition.(15) Inspection of oroantral communications(OAC), especially after maxillary molar and premolar tooth extraction or endodontic surgery performed on maxillary teeth should be done by the surgeon extremely carefully as it can result in sinus perforation which may develop into oroantral communication.(4) The incidence of OAC should be higher after 30 years of life because the maxillary sinus reaches its greatest size during the third decade of life.(6) The incidence rate of 0.31% to 5.1% is there for oroantral communication & subsequent OAF after extraction of upper posterior teeth.(3) OAF occurs after dental extraction. Oroantral fistula is may be the result of several different pathologic processes like infection, inflammatory conditions, neoplasm, Paget’s disease, iatrogenic injury and trauma.(14) Any communication between the oral cavity and the maxillary sinus which lasts for more than 21 days should be closed surgically in order to avoid further medical problems as per the literature.(8) Different parameters must be taken into account including location and size of defect as well as

Conclusion In this reported case, the “OBTURATOR” to close the oroantral communication (OAC) proved successful. The operated case had no complications and there is no recurrgitation of intra orally the oroantral communication was cured.

REFERENCES

1. Hemant Batra, Govind Jindal Supreet Kaur. Evaluation of different treatment modalities for closure of oro-antral communications and formulation of a rational approach. *J Maxillofac Oral Surg* 9(1):13-18.
2. Wael Mohamed Said Ahmed. Closure of Oroantral Fistula Using Titanium Plate with Transalveolar Wiring. *J. Maxillofac. Oral Surg.* (Jan-Mar 2015)14(1):121-125.
3. Yunus Feyyat Sahin, Togay Muderris, Sami Bercin, Ergun Sevil, Muzaffer Kiris. Chronic Maxillary Sinusitis Associated with an unusual Foreign Body: A Case Report. *Case Reports in Otolaryngology*. Volume 2012, Article ID 903714,4 pages.
4. Kiran Kumar Krishanappa S, Prashanti E, Sumanth KN, Naresh S, Moe S, Aggarwal H, Mathew RJ. Interventions for treating oro-antral communications and fistula due to dental procedures (protocol). 2015 The Conchrane Collaboration. Published by John Wiley & sons, Ltd.
5. JosueHernando,LorenaGallego,LuisJunquera, Pedro Villarreal. Oroantral Communications. A retrospective analysis. *Med Oral Patol Oral Cir Bucal*.2010 May 1;15(3):e499-503. 7. Pierre P. Pourmand, Guido R. Sigron, Beatrice Mache, Bernd Standlinger, Michael C. L
6. Andrea Enrico Borgonovo, Frederick Valerio Berardinelli, Marco Favale and Carlo Maiorana-Surgical Options in Oroantral Fistula Treatment. *The Open Dentistry Journal* ,2012,6,94-98.
7. Adele Scattarella, Andrea Ballini, Felice Roberto Grassi, Andrea Carbonara, Francesco Ciccolella, Angela Dituri, Gianna Maria Nardi, Stefania Cantore, Francesco Pettini. Treatment of oroantral fistula with autologous bone graft and application of non-resorbable membrane. *Int.J.Med.Sci*.2010,7.
8. Minutha R, Sriram Nathan.”A study of Caldwell-luc approach in various etiologies”. *Journal of Evolution of Medical and Dental Sciences* 2013;Vol 2,Issue 36,September 9;page:6715-6723.
9. Pedro Henrique de Souza, Lopes, Diogo de Oliveira Sampaio, and Belmino Carlos Amaral Torres. Combined Palatal flap ant titanium mesh for oroantral fistula closure. *Ann Maxillofac Surg*.2015 Jan- Jun;5(1):89-92.
10. Marta del Rey Santamaria, Eduard Valmaseda Castellon, Leonardo Berini Aytes, Cosme Gay Escoda. Incidence of oral sinus communications in 389 upper third molar extraction. *Med Oral Patol Oral Cir Bucal* 2006;11:E334- 8.
11. James J Abrahams, Scott B. Berger. Oral-Maxillary Sinus Fistula (Oroantral fistula): Clinical Features and Findings on Multiplanar CT. *AJR* 1995;165:1273-1276.
12. Fahimeh Akhlaghi, Mohammad Esmaeelinejad, and Pooria Safai. Etiologies and Treatments of Odontogenic Maxillary Sinusitis: A Systematic Review. *Iran Red Crescent Med J*.2015 December;17(12):e25536